THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 18, 1986

EDITORS

M. L. McGlashan
Department of Chemistry
University College London
20 Gordon Street
London WC1H 0AJ
U.K.

P. A. G. O'Hare
Argonne National Laboratory
Chemical Technology Division
Argonne
Illinois 60439
U.S.A.

Assistant Editor: **M. B. Ewing** (University College London, U.K.)

ADVISORY BOARD

Chairman: G. M. Schneider (F.R.G.)

S. Angus (U.K.)
G. C. Benson (Canada)
J. C. G. Calado (Portugal)
M. Colomina (Spain)
J. D. Cox (U.K.)
M. Gaune-Escard (France)
C. Jolicoeur (Canada)
O. J. Kleppa (U.S.A.)
M. Månsson (Sweden)
V. A. Medvedev (U.S.S.R.)
J. B. Ott (U.S.A.)
F. Pavese (Italy)
G. Somsen (The Netherlands)
H. Suga (Japan)
F. L. Swinton (U.K.)
A. G. Williamson (N.Z.)
R. H. Wood (U.S.A.)
W. A. Wóycicki (Poland)

ACADEMIC PRESS
London Orlando San Diego
San Francisco New York Toronto
Montreal Sydney Tokyo São Paulo
A Subsidiary of Harcourt Brace Jovanovich, Publishers

Copyright © 1986 Academic Press Inc. (London) Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 18

Number 1, January 1986

OTT, J. B.; STOUFFER, C. E.; CORNETT, G. V.; WOODFIELD, B. F.; WIRTHLIN, R. C.; CHRISTENSEN, J. J.; DEITERS, U. K. Excess enthalpies for (ethanol + water) at 298.15 K and pressures of 0.4, 5, 10, and 15 MPa . . 1

LEPORI, L.; MATTEOLI, E. Excess volumes of (tetrachloromethane + an alkanol or + a cyclic ether) at 298.15 K 13

LEGIDO, J. L.; BRAVO, R.; PAZ ANDRADE, M. I.; ROMANI, L.; SARMIENTO, F.; ORTEGA, J. Excess enthalpies of five examples of (2-hexanone + an *n*-alkane) and five of (2-hexanone + an *n*-alkanol) at 298.15 K . . 21

KUMARAN, M. K.; BENSON, G. C. Excess enthalpies of (diether + *n*-alkane) at 298.15 K 27

GUANQUAN, C.; OTT, J. B.; GOATES, J. R. (Solid + liquid) phase equilibria and solid-compound formation in 1,2-dimethoxyethane + tetrachloromethane, + trichlorofluoromethane, and + trichloromethane . . . 31

NAGATA, I.; TAMURA, K. Excess molar enthalpies of chlorobenzene + methanol, + acetonitrile, and + ethanol, and of (acetonitrile + chlorobenzene) + methanol and + ethanol at 298.15 K . . . 39

AMBROSE, D. The correlation and estimation of vapour pressures. IV. Observations on Wagner's method of fitting equations to vapour pressures 45

CHRISTENSEN, J. J.; CORDRAY, D.; IZATT, R. M. The excess enthalpies of (carbon dioxide + decane) from 293.15 to 573.15 K at 12.50 MPa . 53

TANAKA, R.; TOYAMA, S.; MURAKAMI, S. Heat capacities of $\{xC_nH_{2n+1}OH + (1-x)C_7H_{16}\}$ for $n = 1$ to 6 at 298.15 K . . . 63

JACKSON, P. R.; MORCOM, K. W. (Solid + liquid) phase equilibria for mixtures containing hexafluorobenzene 75

NAUMANN, D.; WAGNER, H. GG. Vapour-liquid equilibria of several mixtures containing 2-butanone, 2-butenal, ethoxyethanol, and toluene at 330.15 K 81

CHHOR, K.; BOCQUET, J. F.; POMMIER, C.; CHARDON, B. Heat capacity and thermodynamic behaviour of Mn_3O_4 and $ZnMn_2O_4$ at low temperatures 89

Number 2, February 1986

DEWAN, R. K.; MEHTA, S. K. Excess volumes of *n*-octanol + acetonitrile, + butyronitrile, + nitromethane, and + nitroethane 101

SNOW, R. L.; OTT, J. B.; GOATES, J. R.; MARSH, K. N.; O'SHEA, S.; STOKES, R. H. (Solid + liquid) and (vapor + liquid) phase equilibria and excess enthalpies for (benzene + *n*-tetradecane), (benzene + *n*-hexadecane), (cyclohexane + *n*-tetradecane), and (cyclohexane + *n*-hexadecane) at 293.15, 298.15, and 308.15 K. Comparison of G_m^E calculated from (vapor + liquid) and (solid + liquid) equilibria 107

AWWAD, A. M.; AL-MASHHADANI, A. M.; TAWFEEK, E. A. Excess volumes of (*N*-methylpyrrolidone + isoalkanol) at 298.15 K 131

WORMALD, C. J.; LANCASTER, N. M.; SELLARS, A. J. The excess molar enthalpies of $\{xH_2O+(1-x)CO\}$(g) and $\{xH_2O+(1-x)CO_2\}$(g) at high temperatures and pressures 135

ZHANG, D.; BENSON, G. C.; KUMARAN, M. K.; LU, B. C.-Y. Excess thermodynamic properties for methyl methacrylate + methanol and + ethanol 149

MEISINGSET, K. K.; GRØNVOLD, F. Thermodynamic properties and phase transitions of salt hydrates between 270 and 400 K. IV. $CaCl_2 \cdot 6H_2O$, $CaCl_2 \cdot 4H_2O$, $CaCl_2 \cdot 2H_2O$, and $FeCl_3 \cdot 6H_2O$ 159

WAKABAYASHI, T.; TAKAIZUMI, K.; TANAKA, R. Heat capacities of quaternary phosphonium chlorides and of mixed salts in aqueous solutions at 298.15 K 175

O'HARE, P. A. G.; JOHNSON, G. K. Thermochemistry of inorganic sulfur compounds. VII. Standard molar enthalpy of formation at 298.15 K, high-temperature enthalpy increments, and other thermodynamic properties to 1100 K of titanium disulfide, TiS_2 189

Number 3, March 1986

APELBLAT, A.; TAMIR, A. Enthalpy of solution of lithium bromide, lithium bromide monohydrate, and lithium bromide dihydrate, in water at 298.15 K 201

TRESZCZANOWICZ, T.; LU, B. C.-Y. Isothermal vapour-liquid equilibria for 11 examples of (an ether + a hydrocarbon) 213

GEE, N.; SHINSAKA, K.; DODELET, J.-P.; FREEMAN, G. R. Dielectric constant against temperature for 43 liquids 221

KOROBOV, M. V.; NIKULIN, V. V.; CHILINGAROV, N. S.; SIDOROV, L. N. Enthalpy of formation of platinum hexafluoride 235

SHINMEI, M.; IMAI, T.; YOKOKAWA, T.; MASSON, C. R. Thermodynamic study of $Si_2OF_6(g)$ from 723 to 1288 K by mass spectrometry 241

SCHWITZGEBEL, G.; FLOHR, G.; ZORNER, S. E.m.f. and calorimetric investigations on lead germanates 247

AZAD, A. M.; PANKAJAVALLI, R.; SREEDHARAN, O. M. Thermodynamic stability of Sb_2O_3 by a solid-oxide electrolyte e.m.f. technique . . 255

HOLMES, H. F.; MESMER, R. E. Isopiestic studies of aqueous solutions at elevated temperatures. VIII. The alkali-metal sulfates 263

OTSUKA, S.; KOZUKA, Z. The activity coefficient of oxygen in liquid (germanium + tin) and (germanium + antimony) alloys 277

KANA'AN, A. S.; GONZALEZ, C.; KANAMUELLER, J. K. Equilibrium decomposition pressures of thallium(I) hexafluorophosphate 285

RIBEIRO DA SILVA, M. D. M. C.; RIBEIRO DA SILVA, M. A. V.; PILCHER, G. Enthalpies of combustion of the three trihydroxybenzenes and of 3-methoxycatechol and 4-nitrocatechol 295

NOTE

AIROLDI, C.; SANTOS, M. R. M. C. Calcium–oxygen bond enthalpy in hexakis(urea) calcium bromide 301

Number 4, April 1986

RODRIGUEZ-NUÑEZ, E.; PAZ-ANDRADE, M. I.; ORTEGA, J. Thermodynamic properties of (a xylene + an *n*-alkanol). IV. Excess molar volumes at 298.15 K for 1-propanol, 1-butanol, and 1-pentanol 303

MAIER, J.; WARHUS, U. Thermodynamic investigations of Na_2ZrO_3 by electrochemical means 309

WÓYCICKI, W. Excess enthalpies of (cyclohexylamine or aniline + cyclohexane or cyclohexene or a cyclohexadiene) 317

GUANQUAN, C.; OTT, J. B.; GOATES, J. R. (Solid + liquid) phase equilibria and solid-compound formation in tetrachloromethane + cyclopentanone, + 4-butyrolactone, and + 1-methyl-2-pyrrolidinone 323

HNĚDKOVSKÝ, L.; CIBULKA, I. Excess molar volumes of binary liquid mixtures of {an oxygenated compound, or cyclopentane, or pentane + 1-bromo-1-chloro-2,2,2-trifluoroethane (halothane)} at 298.15 K 331

TANIEWSKA-OSIŃSKA, S.; JÓŹWIAK, M. Calorimetric investigations of solutions of NaI, KI, NaCl, KCl, and $C_6H_5NH_2$ in $[xPO\{N(CH_3)_2\}_3 + (1-x)H_2O]$ at 298.15 K 339

APELBLAT, A. Enthalpy of solution of oxalic, succinic, adipic, maleic, malic, tartaric, and citric acids, oxalic acid dihydrate, and citric acid monohydrate in water at 298.15 K 351

STOKES, R. H. Enthalpies of mixing by mixture-displacement calorimetry . 359

BEYER, R. P.; FERRANTE, M. J. Heat capacity of cadmium oxysulfate from 5.16 to 300.7 K and relative enthalpy to 1001.5 K 365

YERLETT, T. K.; WORMALD, C. J. The enthalpy of acetone 371

GRØNVOLD, F.; WESTRUM, E. F., JR. Silver(I) sulfide: Ag_2S. Heat capacity from 5 to 1000 K, thermodynamic properties, and transitions . . 381

Number 5, May 1986

ORTEGA, J.; BRAVO, R.; JIMÉNEZ, E.; PAZ ANDRADE, M. I. Excess molar volumes of (ethyl formate + each of several alkanols) at 298.15 K . 403

SAITO, K.; ATAKE, T.; CHIHARA, H. Thermodynamic studies of *p*-polyphenyls: heat capacity of 4,4′-difluorobiphenyl 407

GOODMAN, M. A.; JAIN, V. K.; WHITTENBURG, S. L. Hypersonic speed and isentropic compressibility in *m*- and *p*-cresols 415

ORTEGA, J.; MATOS, S.; PAZ ANDRADE, M. I.; FERNANDEZ, J.; ROMANI, L. Excess volumes of (toluene + each of several isomers of hexanol) at 298.15 K 419

AMITIN, E. B.; MINENKOV, YU. F.; NABUTOVSKAYA, O. A.; PAUKOV, I. E. Thermodynamic properties of solid mercury from 150 to 234 K . . 423

HALLÉN, D.; NILSSON, S.-O.; ROTHSCHILD, W.; WADSÖ, I. Enthalpies and heat capacities for *n*-alkan-1-ols in H_2O and D_2O 429

AWWAD, A. M. Excess molar volumes for (2-ethoxyethanol + a glycol) at 298.15 K 443

KIMURA, T.; TAKAGI, S. Thermodynamics of liquid mixtures containing methyl methylthiomethyl sulfoxide. I. Excess volumes of (water or benzene or dimethyl sulfoxide + methyl methylthiomethyl sulfoxide) and (water or benzene + dimethyl sulfoxide) at 298.15 and 318.15 K 447

BANNISTER, M. J. Oxygen concentration-cell determination of the standard molar Gibbs free energy of formation of SnO_2 455

O'NEILL, H. ST. C. Standard molar Gibbs free energies of formation of the tetragonal and hexagonal forms of germanium dioxide 465

GRIMSEY, E. J.; REYNOLDS, K. A. Equilibrium oxygen pressure of $(Co_3O_4 + CoO)$ from 1173 to 1228 K 473

STAVELEY, L. A. K.; WEIR, R. D. The heat capacity of paraperiodic acid H_5IO_6 from 8.0 to 305 K 477

JACKOWSKI, A. W. Vapour pressure of (*n*-hexane + *n*-butyl octadecanoate). II. (Solid + liquid + vapour) equilibria 483

TAMURA, K.; MURAKAMI, S.; DOI, S.; KIMURA, T.; TAKAGI, S. Speeds of sound, densities, and isentropic compressibilities for $x \ll 1$ of $\{xo\text{-}C_6H_4(C_6H_5)_2 + (1-x)C_6H_6\}$ and $\{xp\text{-}C_6H_4(C_6H_5)_2 + (1-x)C_6H_6\}$ at 298.15 K . 489

BENSON, G. C.; D'ARCY, P. J. Excess isobaric heat capacities of some binary mixtures: (a C_5-alkanol + *n*-heptane) at 298.15 K 493

NOTES

AUFFRÉDIC, J.-P.; MARTINEZ, O. G.; TIGERAS, P.; DELGADO, A. L. Standard enthalpies of solution and formation of $\beta\text{-}Co_2(OH)_3Cl$ 499

CORDFUNKE, E. H. P.; PRINS, G. The standard enthalpy of formation of UP_2O_7; a correction 501

Number 6, June 1986

CORDFUNKE, E. H. P.; OUWELTJES, W.; PRINS, G. Standard enthalpies of formation of uranium compounds. XIII. Cs_2UO_4 503

EWING, M. B.; MCGLASHAN, M. L.; TRUSLER, J. P. M. The speed of sound in gases. II. Acoustic virial coefficients and perfect-gas heat capacities for 2,2-dimethylpropane obtained using a cylindrical interferometer . . 511

WHITE, M. A.; WAGNER, B. D. A calorimetric investigation of polymorphism in a layered perovskite: $KAlF_4$ 519

OKADA, M.; UEMATSU, M.; WATANABE, K. Orthobaric liquid densities of trichlorofluoromethane, dichlorodifluoromethane, chlorodifluoromethane, 1,1,2-trichlorotrifluoroethane, 1,2-dichlorotetrafluoroethane, and of the azeotropic mixture of (chlorodifluoromethane + chloropentafluoroethane) between 203 and 463 K 527

LANCASTER, N. M.; WORMALD, C. J. Excess molar enthalpies for $\{xH_2O+(1-x)C_3H_6\}$(g), $\{xH_2O+(1-x)C_3H_8\}$(g), and $\{xH_2O+(1-x)C_4H_{10}\}$(g) 545

GARCIA, M.; REY, C.; PEREZ VILLAR, V.; RODRIGUEZ, J. R. Excess volumes of (*n*-heptane + *n*-undecane) between 288.15 and 308.15 K 551

O'HARE, P. A. G. The standard molar enthalpy of formation at 298.15 K of crystalline germanium diselenide ($GeSe_2$). Thermochemical Ge–Se bond-dissociation enthalpies. Revised values for the standard molar enthalpies of formation of GeF_4(g) and SeF_6(g) 555

FLANIGAN, D. A.; YESAVAGE, V. F.; KIDNAY, A. J. Enthalpy of *m*-cresol between 291 and 655 K and at pressures of 207 and 690 kPa . . . 563

OTIN, S.; FERNANDEZ, J.; VEAMONTE, M.; VELASCO, I.; GUTIÉRREZ LOSA, C. Excess molar enthalpies of mixtures containing halogeno-aromatic compounds 571

NUÑEZ, L.; BARRAL, L.; GAVILANES LARGO, S.; PILCHER, G. Enthalpies of combustion of the three aminophenols 575

CORDRAY, D. R.; CHRISTENSEN, J. J.; IZATT, R. M. The excess enthalpies of (propane + chlorotrifluoromethane) from 293.15 to 383.15 K at 4.45 MPa 581

HAMAM, S. E. M.; BENSON, G. C. Excess enthalpies of some binary mixtures of hexane isomers 591

GIDLEY, M. A.; STUBLEY, D. Activity coefficients at infinite dilution of solutes in liquid-crystalline solvents 595

NOTE

ORTEGA, J.; PAZ ANDRADE, M. I.; SARMIENTO, F. Excess molar enthalpies at 298.15 K of (butan-1-ol + ethyl butyl ether) and of (butan-1-ol + ethyl *t*-butyl ether) 601

Number 7, July 1986

GUANQUAN, C.; OTT, J. B.; GOATES, J. R. (Solid + liquid) phase equilibria and solid-compound formation in tetrachloromethane + furan, + pyridine, and + *N*-methylpyrrole 603

NGEYI, S. P.; WESTRUM, E. F., JR.; FRANZOSINI, P. Thermophysics of metal alkanoates. V. Heat capacities and thermodynamic properties of lithium *n*-butanoate and *n*-hexanoate 609

ZHANG, D.; BENSON, G. C.; LU, B. C.-Y. Excess volumes of (*n*-decane + a hexane isomer) at 298.15 K 619

SANAHUJA, A.; GÓMEZ-ESTÉVEZ, J. L. Determination of the enthalpies of solution at infinite dilution of KCl and NaCl in water at 303.15, 308.15, and 313.15 K 623

CÁCERES ALONSO, P.; CÁCERES ALONSO, M.; NÚÑEZ DELGADO, J. Excess volumes of (1,3,5-trimethylbenzene + an *n*-alkane) 629

KAPOSI, O.; BENCZE, L.; ZHURAVLEVA, L. V. Determination of thermodynamic activities by mass spectrometry in $\{xNaI+(1-x)CsI\}$ 635

CORDRAY, D. R.; CHRISTENSEN, J. J.; IZATT, R. M. The excess enthalpies of (carbon dioxide + toluene) at 308.15, 358.15, and 573.15 K from 6.98 to 16.63 MPa 647

NAKANISHI, K.; TOUHARA, H. Excess molar enthalpies of (methanol + aniline), (methanol + *N*-methylaniline), and (methanol + *N*,*N*-dimethylaniline) 657

SIDDIQI, M. A.; LUCAS, K. The excess enthalpy of (trichloroethene + carbon disulphide), (trichloroethene + trichloromethane), and (trichloroethene + tetrachloromethane) at various temperatures and pressures . . 661

CHAUHDRY, M. S.; LAMB, J. A. Excess volumes of (2-butanol + water) at pressures up to 220 MPa 665

NILSSON, S.-O.; WADSÖ, I. Thermodynamic properties of some mono-, di-, and tri-esters. Enthalpies of solution in water at 288.15 to 318.15 K and enthalpies of vaporization and heat capacities at 298.15 K 673

TARDAJOS, G.; DÍAZ PEÑA, M.; AICART, E. Speed of sound in pure liquids by a pulse-echo-overlap method 683

O'SHEA, S. J.; STOKES, R. H. Activity coefficients and excess partial molar enthalpies for (ethanol + hexane) from 283 to 318 K 691

DEWAN, R. K.; MEHTA, S. K. Excess molar volumes of $\{x\underline{CH_2(CH_2)_3}SO_2 + (1-x)C_nH_{2n+1}OH\}$ for $n = 6$, 7, 8, and 10 697

Number 8, August 1986

WIŚNIEWSKA, B.; ŁENCKA, M.; ROGALSKI, M. Vapour pressures of 2,4-, 2,6-, and 3,5-dimethylpyridine at temperatures from 267 to 360 K . . 703

QUEIROZ, J. C.; AIROLDI, C.; CHAGAS, A. P. A thermochemical study involving adducts $MX_2 \cdot P(C_6H_5)_3O$ (M = Zn, Cd, Hg; X = Br, I) . . . 709

KALINOWSKA, B.; JACKOWSKI, A. W. Enthalpies of two phase transitions in the high-temperature form of *n*-butyl octadecanoate 715

YERLETT, T. K.; WORMALD, C. J. The enthalpy of methanol . . . 719

MALLIKA, C.; SREEDHARAN, O. M. Thermodynamic stabilities of TeO_2 and Sb_2Te_3 by a solid-oxide electrolyte e.m.f. technique 727

VENUGOPAL, V.; SHUKLA, N. K.; SUNDARESH, V.; ROY, K. N.; PRASAD, R.; SOOD, D. D. Thermochemistry of caesium iodide and caesium chromate 735

KLEINRAHM, R.; WAGNER, W. Measurement and correlation of the equilibrium liquid and vapour densities and the vapour pressure along the coexistence curve of methane 739

KOKKONEN, P. Excess enthalpies of (triethylamine + *n*-alkanone) at 303.15 and 313.15 K 761

O'HARE, P. A. G.; JENSEN, K. J.; JOHNSON, G. K. Thermochemistry of inorganic sulfur compounds. VIII. Na_2SO_3 and K_2SO_3: standard molar enthalpies of solution and formation at 298.15 K, high-temperature enthalpy increments and heat capacities, and temperatures and enthalpies of transition. A new value for the standard molar enthalpy of formation of SO_3^{2-}(aq) 765

BROWN, R. J. C.; CALLANAN, J. E.; WEIR, R. D.; WESTRUM, E. F., JR. The thermodynamics of ammonium scheelites. II. Heat capacity of deuterated ammonium perrhenate ND_4ReO_4 from 7.5 to 320 K 787

KIRCHNER, J. J.; ACREE, W. E., JR.; PILCHER, G.; SHAOFENG, L. Enthalpies of combustion of four *N*-phenylmethylene benzenamine *N*-oxide derivatives, of *N*-phenylmethylene benzenamine, and of *trans*-diphenyldiazene *N*-oxide: the dissociation enthalpy of the (N–O) bonds 793

NOTE

JOHNSON, G. K. The standard molar enthalpy of formation of SiF_4(g) at 298.15 K by fluorine bomb calorimetry 801

Number 9, September 1986

ROMANI, L.; ARIAS, J. M.; BRAVO, R.; JIMENEZ, E.; PAZ ANDRADE, M. I. Excess molar enthalpies of {an isomer of hexanol + (hexane or cyclohexane or benzene)} at 298.15 K 803

KARVO, M. Ultrasonic speeds and isentropic compressibilities of sulfolane + benzene, + toluene, + *p*-xylene, and + mesitylene at 303.15 K . . 809

OTT, J. B.; HÖLSCHER, I. F.; SCHNEIDER, G. M. (Liquid + liquid) phase equilibria in (methanol + heptane) and (methanol + octane) at pressures from 0.1 to 150 MPa 815

NAGATA, I.; TAMURA, K. Excess molar enthalpies of {tetrachloromethane + propan-1-ol or + propan-2-ol} and of {(tetrachloromethane + benzene) + propan-1-ol or + propan-2-ol} at 298.15 K 827

BERGMAN, C.; CHASTEL, R.; MATHIEU, J.-C. Determination of the thermodynamic properties of (copper + silicon) melts by Knudsen-cell mass spectrometry 835

MAGEE, J. W.; KOBAYASHI, R. Isochoric (p, ρ_n, T) measurements on ($0.2005H_2$ + $0.7995CH_4$) at temperatures from 140 to 273.15 K and pressures to 70 MPa 847

KARAKAYA, I.; THOMPSON, W. T. Thermodynamic properties of $\{(1-x)MgCl_2+xCaCl_2\}(l)$ 859

OTT, J. B.; CORNETT, G. V.; STOUFFER, C. E.; WOODFIELD, B. F.; GUANQUAN, C.; CHRISTENSEN, J. J. Excess enthalpies of (ethanol + water) at 323.15, 333.15, 348.15, and 373.15 K and from 0.4 to 15 MPa 867

NILSSON, S.-O. Enthalpies of solution of water in benzene and in some *n*-alkanes 877

GAMBOA, I.; TARDAJOS, G.; DÍAZ PEÑA, M.; AICART, E. Isothermal compressibility of (toluene + *n*-decane) and (toluene + *n*-dodecane) at various temperatures 885

HANDA, Y. P. Calorimetric determinations of the compositions, enthalpies of dissociation, and heat capacities in the range 85 to 270 K for clathrate hydrates of xenon and krypton 891

Number 10, October 1986

CASQUERO RUIZ, J. DE D.; GUIL PINTO, J. M.; PÉREZ MASIÁ, A.; RUIZ PANIEGO, A. Thermodynamics of the adsorption of hydrocarbons on alumina 903

HANDA, Y. P. Compositions, enthalpies of dissociation, and heat capacities in the range 85 to 270 K for clathrate hydrates of methane, ethane, and propane, and enthalpy of dissociation of isobutane hydrate, as determined by a heat-flow calorimeter 915

PRICE, J. S.; TASKER, I. R.; APPELMAN, E. H.; O'HARE, P. A. G. Thermochemistry of inorganic sulfur compounds. IX. Molar heat capacity of $KHSO_5$(cr) from 5 to 300 K, and the partial molar entropy of HSO_5^-(aq) 923

CALADO, J. C. G.; REBELO, L. P. N.; STREETT, W. B.; ZOLLWEG, J. A. Thermodynamics of liquid (dimethylether + xenon) 931

YAMAMURO, O.; OGUNI, M.; MATSUO, T.; SUGA, H. Calorimetric and dilatometric studies on the phase transitions of crystalline CH_3NH_3I . 939

BERNAL, P. J.; VAN HOOK, W. A. Apparent molar volumes, isobaric expansion coefficients, and isentropic compressibilities, and their H/D isotope effects for some aqueous carbohydrate solutions. 955

BERNAL, P. J.; VAN HOOK, W. A. Apparent molar volumes, isobaric expansion coefficients, and isentropic compressibilities for some non-aqueous carbohydrate solutions 969

EWING, M. B.; JOHNSON, K. A.; MCGLASHAN, M. L. The liquid–liquid critical state of (cyclohexane + methanol). II. Use of the parametric equation of state to represent excess enthalpy in the critical region 979

KUMARAN, M. K.; BENSON, G. C. Excess enthalpies of *n*-dodecane + *n*-heptane, +*n*-octane, and +*n*-decane at 298.15 K 993

KOWALSKA, M.; SKUDLARSKI, K.; BOTOR, J. The thermodynamic activity of SiO_2 in $\{x_1Cu_2O + x_2K_2O + (1-x_1-x_2)SiO_2\}$(l) at 1373 K . . 997

Number 11, November 1986

LOPEZ, M.; PAZ-ANDRADE, M. I.; FERNANDEZ, J.; RODRIGUEZ-NUÑEZ, E.; ORTEGA, J. Excess molar enthalpies at 298.15 K of (an *n*-alkyl formate + an *n*-alkanol). I. $\{xHCO_2(CH_2)_3CH_3+(1-x)C_nH_{2n+1}OH\}$, ($n = 3$ to 10) 1003

BENSON, G. C.; KUMARAN, M. K.; HALPIN, C. J.; D'ARCY, P. J. Thermodynamic properties for (3,3-dimethylbutan-1-ol + *n*-hexane) at 298.15 K 1007

DEWAN, R. K.; MEHTA, S. K. Excess molar volumes of ethylbenzene + acetonitrile, + butyronitrile, + nitromethane, and + nitroethane . . 1015

SINGH, R. R.; VAN HOOK, W. A. Effect of dissolved gas on the critical solution temperature of (cyclohexane + methanol) 1021

JUNEJA, J. M.; IYENGAR, G. N. K.; ABRAHAM, K. P. Thermodynamic properties of liquid (magnesium + copper) alloys by vapour-pressure measurements made by a boiling-temperature method 1025

LETCHER, T. M.; WOOTTON, S.; SHUTTLEWORTH, B.; HEYWARD, C. Phase equilibria for (*n*-heptane + water + an alcohol) at 298.2 K . . 1037

LUO, B.; HAMAM, S. E. M.; BENSON, G. C.; LU, B. C.-Y. Excess enthalpies and excess volumes of (methyl methacrylate + *n*-hexane) and (methyl methacrylate + *n*-heptane) 1043

NAUMOWICZ-WĘGLIŃSKA, E.; WÓYCICKI, W. Excess enthalpies of gaseous mixtures containing ammonia. 2. (Ammonia + nitrogen) and (ammonia + methane) 1047

CHRISTENSEN, J. J.; FAUX, P. W.; CORDRAY, D.; IZATT, R. M. The excess enthalpies of (carbon dioxide + pentane) at 348.15, 373.15, 413.15, 470.15, and 573.15 K from 7.58 to 12.45 MPa 1053

MATTEOLI, E.; LEPORI, L. Excess Gibbs free energies of (tetrachloromethane + an alcohol or a cyclic ether) at 298.15 K 1065

HENDERSON, C.; LEWIS, D. G.; PRICHARD, P. C.; STAVELEY, L. A. K.; FONSECA, I. M. A.; LOBO, L. Q. Some thermodynamic properties of hydrogen chloride and deuterium chloride 1077

PALACIOS, E.; NAVARRO, R.; BURRIEL, R.; BARTOLOMÉ, J.; GONZÁLEZ, D. Thermal properties of XMF_3: cubic perovskites. I. Heat capacity of NH_4MgF_3 and NH_4CdF_3 1089

Number 12, December 1986

KLEINRAHM, R.; DUSCHEK, W.; WAGNER, W. (Pressure, density, temperature) measurements in the critical region of methane 1103

NILSSON, S.-O. Partial molar enthalpies of solution of H_2O in some *n*-alkan-1-ols and esters at 298.15 and 308.15 K 1115

NILSSON, S.-O.; WADSÖ, I. A flow-microcalorimetric vessel for solution of slightly soluble solids 1125

NAVARRO, R.; BURRIEL, R.; BARTOLOMÉ, J.; GONZÁLEZ, D. Thermal properties of XMF_3 cubic perovskites. II. Heat capacity of NH_4ZnF_3 and $KZnF_3$ 1135

BENSON, G. C.; HALPIN, C. J.; KUMARAN, M. K. Ultrasonic speeds and isentropic compressibilities of each of the mixtures: (*n*-dodecane + an isomer of hexane) at 298.15 K 1147

ROSENHOLM, J. B.; GRIGG, R. B.; HEPLER, L. G. Thermodynamic properties of aqueous solutions of surfactants: molar heat capacities and volumes 1153

KOKKONEN, P. Excess molar enthalpies of triethylamine + ethylbenzene, + *n*-propylbenzene, + *n*-butylbenzene, + isopropylbenzene, and + isobutylbenzene at 303.15 K 1165

FRANZOSINI, P.; NGEYI, S. P.; WESTRUM, E. F., JR. Thermophysics of metal alkanoates. VI. Heat capacities and thermodynamic properties of lithium *n*-pentanoate and *n*-heptanoate 1169

SMITH, M. J.; VINCENT, C. A. Determination of thermodynamic formation values for some (silver + transition-metal dichalcogenide) intercalation compounds 1183

AUFFREDIC, J. P.; GARCIA-MARTINEZ, O.; TIGERAS, P.; LOPEZ-DELGADO, A. Standard enthalpies of formation of lanthanum hydroxynitrates . . 1189

MARTIN, J. F.; DOWNIE, D. B. Heat capacities of transition-metal alloys. 3. Sigma-phase alloy $Mn_{0.794}Cr_{0.206}$ 1193

NOTE

El-Hadi, M. F.; Metwally, M. S. Thermodynamic quantities in the exchange of alkali-metal ions on variously cross-linked *m*-cresol formaldehyde resins 1203

Index to Volume 18 1205